

**ENDOGENOUS IRRIGATION:
THE IMPACT OF CLIMATE CHANGE ON FARMERS IN AFRICA**

Pradeep Kurukulasuriya and Robert Mendelsohn¹

¹ School of Forestry and Environmental Studies, Yale University, 230 Prospect St, New Haven, CT 06511, USA. E-mails: pradeep.kurukulasuriya@yale.edu; robert.mendelsohn@yale.edu

The authors would like to thank Rashid Hassan, James Benhin and Glwadys Gbetibouo, Centre for Environmental Economics and Policy in Africa (CEEPA), University of Pretoria, South Africa; Mbaye Diop and Isidor Sène, ISRA/LERG, Campus Universitaire de l'ESP, Senegal; Helmy Mohamed Eid, Samia El-Marsafawy and Samiha Ouda, Soil, Water & Environment Research Institute (SWERI), Egypt; K Yerfi Fosu and Joseph Adu, Department of Agricultural Economics, University of Ghana; Suman Jain Mathematics and Statistics Department, University of Zambia; David Maddison, Department of Economics, University College London, UK; Ali Mahamadou and M Katiella, Faculty of Sciences, University of Abdou Moumouni of Niamey, Niger; Leopold Some and Mathieu Ouedraogo, Institut de l'Environnement et de Recherches Agricoles (INERA), Burkina Faso; and Ariel Dinar, Agricultural and Rural Development Department, World Bank, USA. The project was funded by the Global Environment Facility, the World Bank, and CEEPA.

We are deeply grateful to the country teams from the 11 countries in the study for designing, collecting, and cleaning the data and making this project a success.

All errors are solely the responsibility of the authors and should not be ascribed to any of the institutions or persons listed above.

THE MISSION OF CEEPA

The mission of the Centre for Environmental Economics and Policy in Africa (CEEPA) is to enhance the capacity of African researchers to conduct environmental economics and policy inquiry of relevance to African problems and increase the awareness of the role of environmental economics in sustainable development for economic managers and policy makers.

CEEPA PUBLICATIONS

Aims and Scope

CEEPA publishes peer-reviewed work in environmental economics in its Discussion Paper Series and other publications. The goal of the publications is to promote the exchange of ideas among environmental economists conducting both quantitative and qualitative analysis for both analytical and applied policy design in the public and private sectors in Africa. CEEPA also publishes research materials intended as useful resources for researchers, policy makers and students in environmental economics and policy.

- Co-Editors:** Prof Margaret Chitiga, University of Pretoria
Dr Eric Dada Mungatana, CEEPA
- Associate Editors:** Dr Tim Lynam, Tropical Resources and Ecology Policy Studies (TREPS), University of Zimbabwe
Prof Mohamud Jama, Institute for Development Studies, University of Nairobi
Dr Ayalneh Bogale, Department of Agricultural Economics, Alemaya University
- Assistant Editor:** Dalène du Plessis, CEEPA

Correct citation: Kurukulasuriya P & Mendelsohn R, 2006. Endogenous irrigation: The impact of climate change on farmers in Africa. CEEPA Discussion Paper No. 18, Centre for Environmental Economics and Policy in Africa, University of Pretoria.

Core funding for CEEPA comes from the University of Pretoria. CEEPA receives supplementary funding from various donors including the Swedish International Development Cooperation Agency (SIDA), GEF, NRF South Africa, NOAA (USA), IDRC.

Additional information on CEEPA activities is available at <http://www.ceepa.co.za>
Department of Agricultural Economics, Faculty of Natural and Agricultural Sciences
University of Pretoria; PRETORIA 0002, Republic of South Africa

ISBN 1-920160-18-3
First published 2006
Series ISBN 1-920160-01-9

Printed and bound in South Africa by University of Pretoria

© 2006, Centre for Environmental Economics and Policy in Africa (CEEPA)

TABLE OF CONTENTS

Section	Page
Preface	3
Executive summary	4
1 Introduction	5
2 Model	6
3 Empirical results	7
4 Climate change simulation	10
5 Conclusions	11
References	12
Appendix A: Temperature normals (Sample means)	14
Appendix B: Precipitation normals (Sample means)	14

LIST OF TABLES

Table	Page
1 Sample of farms	15
2 Probit model of whether to irrigate	16
3 Net revenue regressions	18
4a Seasonal and annual marginal effects, evaluated at the sample means for irrigated farms	21
4b Seasonal and annual marginal effects, evaluated at the sample means for dryland farms	22
5 Marginal annual climate impacts	22
6 Comparison of irrigation and welfare estimates for endogenous and exogenous models across different climate scenarios	23

LIST OF FIGURES

Figure	Page
1a Relationship between seasonal temperature and the probability of adopting irrigation	24
1b Relationship between annual temperature and the probability of adopting irrigation	24
2a Relationship between seasonal precipitation and the probability of adopting irrigation	25
2b Relationship between annual precipitation and the probability of adopting irrigation	25
3 Temperature response functions of irrigated and dryland farms	26
4 Precipitation response functions of irrigated and dryland farms	26

PREFACE

PREFACE

The reports in this special series are the result of a multi-country research activities conducted under the GEF funded project: *Climate Change Impacts on and Adaptation of Agro-ecological Systems in Africa*. The main goal of the project was to develop multipliable analytical methods and procedures to assess quantitatively how climate affects current agricultural systems in Africa, predict how these systems may be affected in the future by climate change under various global warming scenarios, and suggest what role adaptation could play. The project has been implemented in 11 countries: Burkina Faso, Cameroon, Ghana, Niger and Senegal in west Africa; Egypt in north Africa; Ethiopia and Kenya in east Africa and South Africa, Zambia, and Zimbabwe in southern Africa. The study countries covered all key agro-climatic zones and farming systems in Africa. This is the first analysis of climate impacts and adaptation in the Africa continent of such scale and the first in the world to combine cross-country, spatially referenced survey and climatic data for conducting this type of analysis.

The analyses reported in this series focus mainly on quantitative assessment of the economic impacts of climate change on agriculture and the farming communities in Africa, based on both the cross-sectional (Ricardian) method and crop response simulation modeling. The cross sectional analysis also allowed for assessing the possible role of adaptation. Moreover, the project employed river-basin hydrology modeling to generate additional climate attributes for the impact assessment and climate scenario analyses such as surface runoff and streamflow for all districts in the study countries.

The Centre for Environmental Economics and policy in Africa (CEEPA) of the University of Pretoria coordinated all project activities in close collaboration with many agencies in the involved countries, the Agriculture and Rural Development (ARD) Department of the World Bank, the World Bank Institute (WBI), the Food and Agriculture Organization (FAO), Yale University, the University of Colorado, and the International Water Management Institute (IWMI). The project received supplemental funding from TFESSD, Finnish TF, NOAA-OPG, and CEEPA. We are grateful for the invaluable contributions of all these institutions and all individuals involved in this project. All opinions presented in this report series and any errors in it are those of the authors and do not represent the opinion of any of the above listed agencies.

Rashid Hassan, Project Leader
CEEPA, University of Pretoria

Ariel Dinar, Project Manager
ARD, World Bank

EXECUTIVE SUMMARY

Previous Ricardian analyses of agriculture have either omitted irrigation or treated irrigation as though it is exogenous. In practice, it is a choice by farmers that is sensitive to climate. This paper develops a choice model of irrigation in the context of a Ricardian model of cropland. We first examine how climate affects the decision to employ irrigation and then how climate affects the net revenues of dryland and irrigated land. This Ricardian ‘selection’ model, using a modified Heckman model, is then estimated across 8400 farmers in Africa. We explicitly model irrigation, but we control for the endogeneity of irrigation that plagues a recently suggested remedy.

We find that the choice of irrigation is sensitive to both temperature and precipitation. Simulating the welfare impacts of several climate scenarios, we demonstrate that a model which assumes irrigation is exogenous provides a biased estimate of the welfare effects of climate change. If dryland and irrigation are to be estimated separately in the Ricardian model, irrigation must be modeled endogenously.

The results also indicate that African agriculture is sensitive to climate change. Many farmers in Africa will experience net revenue losses from warming. We find that the elasticity of net revenue with respect to temperature is -0.82 for dryland farms. That is, a 10% increase in temperature will lead to a loss in net revenues per hectare, on average, of 8.2%. Irrigated farms, on the other hand, are more resilient to temperature change and, on the margin, are likely to realize slight gains in productivity. However, any reduction in precipitation will be especially deleterious to dryland farmers, generally the poorest segment of the agriculture community. Dryland farms are sensitive to precipitation (elasticity of 0.28) whereas precipitation has virtually no effect on the net revenues of irrigated farms. As long as there is sufficient water, irrigation appears to buffer farms from precipitation. This is a consistent result across all the models tested in this paper.

Our results indicate that irrigation is an effective adaptation against loss of rainfall and higher temperatures provided there is sufficient water available. This will be an effective remedy in select regions of Africa with water. However, for many regions there is no available surface water, so that warming scenarios with reduced rainfall are particularly deleterious.

1. Introduction

The Ricardian method for estimating the impacts of climate change on agriculture is a regression of land values (or net revenue) against climate and other exogenous characteristics (Mendelsohn et al. 1994). A consistent criticism that has been leveled at the first Ricardian study is that it did not properly take into account irrigation (Cline 1996; Darwin 1999; Schlenker et al. 2005). Adding a dummy variable for irrigation does not change the results (Mendelsohn & Nordhaus 1999). However, in US samples, dryland and irrigated land do have different climate response functions (Mendelsohn & Dinar 2003; Schlenker et al. 2005). Based on these results, Schlenker et al. (2005) argue that the welfare effects from climate change should be estimated separately for irrigated and dryland farms and added. However, this approach is problematic because it treats irrigation as though it is exogenous. The decision to irrigate is a choice and this choice is influenced by climate (Mendelsohn & Dinar 2003). Further, there may be sample selection bias if we rely on farms that are observed to use dryland or irrigation.

In this paper, we develop a new Ricardian model that examines dryland and irrigated land separately but treats the choice of irrigation as endogenous. A variety of factors influence the decision on whether to irrigate. Surface flows, soil types, and subsidies all play a role in making this choice. But perhaps more importantly to climate analyses, the choice is sensitive to climate. Studies that assume irrigation is exogenous fail to take into account how irrigation will change as climate changes and therefore provide biased estimates of the impact of climate change. Moreover, these analyses of only irrigated farms and only dryland farms rely on self-selected samples, not random samples. Studies that fail to account for this non-randomness in the modeling framework will be biased (Heckman 1979; Lee 1983). In Section 2 we develop a theoretical model that improves on past efforts to model irrigation with the Ricardian approach by explicitly addressing farmer choice and selection bias.

We tested this model empirically using a sample of over 8400 farmers from across 11 African countries. The results reveal that the choice of irrigation is endogenous. Farmers select irrigation rather than dryland to maximize profits. As temperatures warm or precipitation declines, farmers turn to irrigation to keep their farms viable. As long as there is a sufficient flow of water, irrigation is an important adaptation strategy.

We then used this empirical model to examine the welfare impacts of climate change on African agriculture. Using a mild and a severe climate scenario, we examined how irrigation and net revenues will be affected. We compared the results of our model with endogenous irrigation with a model that assumes irrigation is exogenous. We found evidence of selection bias but, more importantly, we found that treating irrigation as though it is exogenous leads to biased welfare estimates. The paper concludes by summarizing the results and discussing some policy implications.

2. Model

The underlying theoretical structure of this model assumes that each farm maximizes profits:

$$\max \Pi = P_i Q^*(X, E) - WX \quad (1)$$

where Π is profit, P_i is output prices, Q^* is output, X are chosen inputs, E is environmental factors such as climate and soils, and W is the price of inputs. In this paper, we assume that the amount of cropland is fixed, in order to focus on the issue of irrigation.²

Formally, we rely on an approach similar to the sample selection model for labor (Heckman 1979). However, there is an important difference. In the labor example, people who did not work had no observed income. In this model, farmers who choose not to irrigate still have observed income from dryland farming.

We assume that a farmer irrigates if irrigation is more profitable than dryland farming. In the first stage, we estimate a dichotomous choice model of irrigation, Y , where $Y=1$ is irrigation (1) and $Y=0$ is dryland farming:

$$Y_i = \beta^1 X + \mu_1 \quad (2)$$

In the second stage, we estimate a conditional profit function for each type of farming based on the available exogenous variables, Z :

$$\Pi_i = \gamma^1 Z^1 + \mu_2 \text{ if } Y = 1 \quad (3)$$

$$\Pi_D = \gamma^D Z^D + \mu_3 \text{ if } Y = 0 \quad (4)$$

where Y_1 is a latent variable explaining the choice of irrigation, Π_I is the net profit of farms that have chosen irrigation, and Π_D is the net profit of farms that have chosen dryland farming, X is a k -vector of regressors, Z^I is an m -vector of regressors for irrigation, Z^D is an m -vector of regressors for dryland, and the error terms U_1 and U_2 and U_1 and U_3 are jointly normally distributed, independently of X and Z , with zero expectations.

² Land uses themselves are influenced by climate and other variables (Mendelsohn et al. 1996). However, this topic is beyond the scope of this paper.

$$\begin{aligned}
& u_1 && \sim && N(0,1) \\
& u_2 \sim N(0, \sigma_2) \\
& u_3 && \sim && N(0, \sigma_3) \\
& \text{corr}(u_1, u_2) = \rho_2 \\
& \text{corr}(u_1, u_3) = \rho_3
\end{aligned}$$

Irrigation is observed only if it is more profitable than dryland farming. Thus, the observed dependent variable Y is:

$$Y=1 \text{ if } \Pi_I > \Pi_D$$

$$Y=0 \text{ if } \Pi_D > \Pi_I$$

When $\rho = 0$, OLS (Ordinary Least Squares) regression provides unbiased estimates, but when $\rho \neq 0$ the OLS estimates are biased. We consequently employ the estimated Mills ratio from the selection model in both the irrigated and dryland conditional regressions in order to control for selection (Dubin & McFadden 1984). We expect the signs on the coefficient of the estimated Mills ratio to be opposite in each regression. With the estimated Mills ratios, the selection model allows us to use information on whether farms irrigate or not to improve the estimates of the parameters in the regression model. That is, the selection model provides consistent, asymptotically efficient estimates for all parameters in the model (Dubin & McFadden 1984).

3. Empirical results

The empirical analysis is based on a household survey conducted of 11 countries across Africa: Burkina Faso, Cameroon, Egypt, Ethiopia, Kenya, Ghana, Niger, Senegal, South Africa, Zambia and Zimbabwe (for more information about the entire study, see Dinar et al. 2006). It was difficult to collect land values in this setting. We consequently relied on measures of net revenue per hectare. Net revenue is defined as gross revenue minus the cost of transport, packaging and marketing, storage, post-harvest losses, hired labor (valued at the median market wage rate), light farm tools (such as files, axes, machetes, etc.), rental on heavy machinery (tractors, ploughs, threshers and others), fertilizer and pesticide. Median district prices from the survey were used for both input and crop prices. Household labor costs are not included as a cost in net revenues because it was not clear what value to assign to wages. We controlled for household labor by using household size as a proxy.

In each country, districts were chosen to get farms across a wide range of climate conditions in that country. In each chosen district, a random but clustered sample of farms was selected. The clustering helped to reduce survey expenses. The number of surveys in each country varied but a total of 9597 surveys was administered. After data cleaning, including removal of farms that did not grow crops, and surveys with field errors and missing information, the

final number of useable surveys was 8463. We conducted the analysis at the plot level of each farm as the dataset was sufficiently detailed to extract and utilize information about whether or not a particular plot (from a set of three) was irrigated or not. Each farm provided plot specific data on whether or not irrigation was used, crop production (including crop type, amount harvested, quantity sold, quantity consumed and amount of sales receipt) and crop costs (fertilizer, pesticide and seed data). Using this data, prices per crop and yields per hectare of farmland and cropland were estimated, as well as plot specific crop revenues and farm level gross and net revenues. Net revenue estimates are at the farm level because the input data, including labor (both hired and household) and machinery, were available only at that unit of measurement. It was not possible to allocate most inputs to specific plots as much of it was applied to several plots at a time. The dataset we used contains 1750 irrigated plots and 9183 dryland plots. The distribution of surveys – irrigated and dryland plots by country – is shown in Table 1.

In this study, we relied on monthly temperature data collected from US Department of Defense satellites (Basist et al. 2001). This set of polar orbiting satellites obtain measurements at a given location on earth at 6am and 6pm every day. The satellites are equipped with sensors that measure surface temperature by detecting microwaves that pass through clouds (Weng & Grody 1998). The monthly precipitation data comes from the Africa Rainfall and Temperature Evaluation System (ARTES) (World Bank 2003). This dataset, created by the National Oceanic and Atmospheric Association's Climate Prediction Center, is based on ground station measurements of precipitation over the period 1948–2001. The average temperatures and precipitation for each country in the sample are shown in Appendices A and B. Note that there is a wide range of climates across the 11 countries in the sample.

It is not possible to use every month of climate in a Ricardian regression because of the high correlation between one month and the next. Consequently we must cluster the monthly data into seasons. However, it is not self-evident how to cluster monthly temperatures into a limited set of seasonal measurements. We explored several ways of defining three-month average seasons, starting with November, December, and January for winter. Comparing the results, we found that defining winter in the northern hemisphere as the average of November, December, and January provided the most robust results for Africa. This assumption in turn implies that the next three months would be spring, the three months after that would be summer, and August, September and October would be fall (in the north). These seasonal definitions were chosen because they provided the best fit with the data and reflected the mid-point for key rainy seasons in the sample. We adjusted for the fact that seasons in the southern and northern hemispheres occur at exactly the opposite months of the year.

Soil data was obtained from FAO (2003). The FAO data provides information about the major and minor soils in each location. Data concerning the hydrology was predicted from a hydrological model for Africa (Strzepek & McCluskey 2006). The model calculated the water flow through each district in the surveyed countries. Data on elevation at the centroid of each district was obtained through GIS manipulation using data from the United States Geological Survey (USGS, 2004). The USGS data is derived from a global digital elevation model with a horizontal grid spacing of 30 arc seconds (approximately one kilometer).

During pre-testing of the survey instrument³, it was determined that some African farmers cultivated at least two plots of land. Subsequently, the survey data collected crop data, including production quantities, amount sold, and sale receipts from crops for the largest single plot of cultivated land (referred to hereafter as the main plot) and all others (referred to as the secondary plot). In the following analysis we therefore contend with two plots.

In the first stage of the analysis, we estimated a probit model of whether to irrigate or not (Table 2). We relied on the 10880 plots (out of a total of 10933) for which we have complete information for the regression. The explanatory variables in the first stage included seasonal climate variables, various soils, and flow (millions of m³). We included only the linear climate variables in the first stage. We tested the inclusion of quadratic climate variables but found the linear model to be more reliable. (Log pseudolikelihood = -2340.59 and r-squared 0.51 versus -2187.4434 and 0.54, respectively, for the quadratic probit model). The Chow test for determining the null hypothesis that the estimated parameters are jointly the same is rejected ($\chi^2(30) = 3967.66$; Prob > $\chi^2 = 0.0000$). The coefficients (which are highly significant) suggest that the probability of adoption of irrigation increases with higher temperatures and precipitation in each season except in spring. The reported standard errors in the paper are based on the Huber-White estimator of variance which are robust against many types of misspecification of the model (Heltberg & Tarp 2002). The annual marginal effects, which are more informative of the decision to irrigate or not, reflected in the probability response functions of choosing irrigation given delta temperature increments (holding all other variables constant), reflect the current irrigation landscape in Africa. The probability of adoption of irrigation increases in regions with lower temperatures (for example Egypt and South Africa), while it decreases in warmer regions. Irrigation in cooler regions is more profitable because it requires less water and the crops are more productive. Similarly, in regions of higher precipitation or available flow, the probability of adopting irrigation decreases. Irrigation is less profitable in wetter locations because the fixed cost of irrigation remains the same but the net increment to production declines.

In the probit model, we controlled for water flow by including the log transformation of a long run average (30 years) of estimated mean flow. The coefficient on this variable is positive and significant. In the selection model, we also controlled for soils. The soil variables reflect the proportion of a district with a particular soil type. The inclusion of certain soils specific to a particular region or district results in the model not being full rank (thereby making the interpretation of the statistical significance of the coefficients unreliable). As a result, we included only those soils that are jointly significant for both irrigated and dryland farms.

We then turned to estimating the second stage model of net revenue conditional on type of farm (Table 3, endogenous columns). We used the coefficients of the probit model to estimate the Mills ratio. Following the standard Heckman model, we included the Mills ratio as an additional explanatory variable to control for self-selection bias in the second stage OLS model (Dubin & McFadden 1984). We examined two sets of second stage OLS models: one for dryland and one for irrigated land. The coefficient on the estimated Mills ratio is significant in the dryland regression and negative as anticipated but not significant in the irrigated model. We tested several control variables in each regression (including gender, education and whether the head of the household was a full time farmer), but dropped them because they were not significant.

³ Available on request from the authors.

A comparison of the OLS coefficients in Table 3 confirms our hypothesis that irrigated and dryland farms are different. The log of size of household has a positive effect on net revenue per hectare for both irrigated and dryland farms. Household size is logged because productivity per worker is expected to fall as households become too large. The coefficient of the log of elevation is negative and significant in the dryland equation but it is not significant in the irrigated model. In addition, our findings lend support to the controversial but often observed inverse relationship between farm size and productivity (Sen 1962). Controlling for labor, machinery and other farm inputs, including irrigation and technology, small farms have higher net revenues per hectare than large farms. In our study, net revenue per hectare may be higher because farmers devote more household labor per hectare on smaller farms. We also included a dummy variable that denotes whether or not a farm has electricity. It is clear that electrified farms outperform farms that do not have electricity in both the irrigated and dryland models. Electrification might directly enhance productivity and earnings or it may simply be a proxy for farms that are closer to markets or more modern.

The second stage regressions give an important insight into the climate sensitivity of farms. The results clearly show that dryland and irrigated farms are both sensitive to climate. Evaluating the marginal impact of temperature and precipitation at the mean climate for the sample reveals many significant seasonal impacts (Table 4a,b). In most seasons (except for winter temperature and winter and spring precipitation), the signs of the coefficients for both types of farms are in the same direction. However, the marginal effects of changes in temperature are not the same across seasons. In spring and fall, the marginal temperature effect is negative whereas in summer and winter it is positive.

These offsetting seasonal effects make annual impacts more ambiguous. The annual marginal impacts of temperature and precipitation are shown in Table 5. The magnitudes of the annual temperature effects for dryland and irrigated farms are different. The resulting elasticity of net revenue with respect to temperature is -0.81 and 0.31 for dryland and irrigated farms respectively. The precipitation results for dryland and irrigated land are also quite different. Dryland farms are sensitive to precipitation (elasticity of 0.28) whereas precipitation has virtually no effect on the net revenues of irrigated farms. As long as there is sufficient water, irrigation appears to buffer farms from insufficient precipitation.

In addition to the second stage regressions in Table 3, we also estimated a pair of regressions that treat irrigation as exogenous (Schlenker et al. 2005). The difference is that the first two columns also include the Mills ratio (i.e., first two columns-endogenous adjust for sample selection bias). Although the Mills ratio coefficient is significant in the dryland regression, it is not significant in the irrigated regression. Further, the climate coefficients in the two dryland and two irrigated regressions are quite similar. Sample selection bias does not appear to be an important problem in this dataset. Figures 1 and 2 plot the resultant response functions from the selection model as well as the second stage conditional models for dryland and irrigated farms by varying temperature and precipitation respectively.

4. Climate change simulation

In this section, we calculate the welfare effect of changing climate. We compare the welfare results from our endogenous modeling approach with the welfare results from the exogenous model of irrigation (Schlenker et al. 2005). Note that with the exogenous model of irrigation,

it is assumed that climate change has no effect on the probability of irrigation. The endogenous model allows this probability to change with the climate scenario.

We examined four simple scenarios to illustrate the importance of modeling irrigation correctly. The scenarios assume a uniform change in either temperature or precipitation across Africa. We examined two temperature changes of 2.5°C and 5.0°C warming and a +20% and a -20% change in precipitation. In Table 6, we present the results of each scenario. First, we demonstrated that the different climate change scenarios change the fraction of farms that are irrigated. Second, we showed that the welfare estimates using a model that addresses endogeneity (referred herein as the ‘endogenous model’) and the model that assumes that irrigation is exogenous when it is not (referred to as the ‘exogenous model’) are quite different. Our endogenous model indicates the overall changes in welfare from a 2.5 and 5 degree increase in temperature are -8% and -14% respectively. The exogenous model overestimates the welfare losses in both cases. Our endogenous model predicts that a 20% decrease in precipitation reduces overall welfare by 21% while a 20% increase in precipitation increases welfare by 18%. The exogenous model underestimates both the damages and benefits of these two scenarios respectively. By failing to take into account how farmers change their irrigation decision as climate changes, the exogenous model leads to biased welfare estimates.

5. Conclusions

This paper provided an improved modeling framework for the Ricardian method in analyzing the effect of irrigation on farm performance. We explicitly modeled irrigation as recommended (Cline 1996; Darwin 1999; Schlenker et al. 2005), but we controlled for the endogeneity of irrigation that plagues a recently suggested remedy (Schlenker et al. 2005). Our results indicate that treating irrigation as exogenous leads to biased welfare estimates from climate change. If dryland and irrigation are to be estimated separately in the Ricardian model, irrigation must be modeled endogenously.

The results also indicate that African agriculture is sensitive to climate change. Many farmers in Africa will experience net revenue losses from warming. Any reduction in precipitation will be especially deleterious to dryland farmers, generally the poorest segment of the agriculture community. Irrigation is an effective adaptation against loss of rainfall and higher temperatures provided there is sufficient water available. This will be an effective remedy in select regions of Africa with water (FAO 1997). However, for many regions, there is no available surface water, so that warming scenarios with reduced rainfall are particularly deleterious. On the other hand, mild warming scenarios with increased rainfall may not be harmful at all.

REFERENCES

- Basist A et al., 2001. Using the Special Sensor Microwave Imager to monitor surface wetness. *Journal of Hydrometeorology* 2: 297–308.
- Cline WR, 1996. The impact of global warming on agriculture: Comment. *American Economic Review* 86: 1309–1312.
- Darwin R, 1999. The impacts of global warming on agriculture: A Ricardian analysis: Comment. *American Economic Review* 89: 1049–1052.
- Dubin JA & McFadden DL, 1984. An econometric analysis of residential electric appliance holdings and consumption. *Econometrica* 52(2): 345–362.
- FAO (Food and Agriculture Organization), 1997. Irrigation potential in Africa: A basin approach. *FAO Land and Water Bulletin*, 4, FAO Land and Water Development Division, Rome.
- FAO (Food and Agriculture Organization), 2003. The digital soil map of the world: Version 3.6 (January), Rome, Italy.
- Heckman JJ, 1979. Sample selection bias as a specification error. *Econometrica* 47: 153–161.
- Heltberg R & Tarp F, 2002. Agricultural supply response and poverty in Mozambique. *Food Policy* 27: 103–124.
- Kurukulasuriya P & Mendelsohn R, 2005. A regional analysis of the impact of climate change on African agriculture, Mimeo, Yale University.
- Lee LF, 1983. Generalized econometric models with selectivity. *Econometrica* 51: 507–512.
- Mendelsohn R & Dinar A, 2003. Climate, water, and agriculture. *Land Economics* 79(3): 328–341.
- Mendelsohn R & Nordhaus W, 1996. The impact of global warming on agriculture: Reply. *American Economic Review* 86: 1312–1315.
- Mendelsohn R & Nordhaus W, 1999. The impact of global warming on agriculture: Reply to Darwin. *American Economic Review* 89: 1053–1055.
- Mendelsohn R, Nordhaus W & Shaw D, 1994. The impact of global warming on agriculture: A Ricardian analysis. *American Economic Review* 84: 753–771.
- Mendelsohn R, Nordhaus W & Shaw D, 1996. Climate impacts on aggregate farm values: Accounting for adaptation. *Agriculture and Forest Meteorology* 80: 55–67.
- Schlenker W, Hanemann M & Fischer A, 2005. Will US agriculture really benefit from global warming? Accounting for irrigation in the hedonic approach. *American Economic Review*, 95(1): 395–406.
- Sen AK, 1962. An aspect of Indian agriculture. *Economics Weekly Annual Number*: 243–66.

- Strzepek K & McCluskey A, 2006. District level hydroclimatic time series and scenario analysis to assess the impacts of climate change on regional water resources and agriculture in Africa. CEEPA Discussion Paper No 13, Centre for Environmental Economics and Policy in Africa, University of Pretoria.
- USGS (US Geological Survey), 2004. Global 30 Arc Second Elevation Data, USGS National Mapping Division, EROS Data Centre. (These data files are downloadable from <http://edcdaac.usgs.gov/gtopo30/gtopo30.asp>)
- Weng F & Grody N, 1998. Physical retrieval of land surface temperature using the Special Sensor Microwave Imager. *Journal of Geophysical Research* 103: 8839–8848.
- World Bank, 2003. Africa rainfall and temperature evaluation system (ARTES). World Bank, Washington DC.

APPENDICES

Appendix A: Temperature normals (Sample means)

country	winter	spring	summer	fall
burkinafaso	23.55	28.34	28.87	24.48
cameroon	19.38	21.38	19.97	18.87
egypt	11.67	13.17	24.11	23.38
ethiopia	18.64	21.53	19.71	18.07
ghana	21.79	24.81	22.63	21.16
kenya	18.75	19.72	18.36	19.12
niger	26.28	30.83	33.91	29.18
senegal	24.54	29.13	31.53	26.67
south africa	11.53	15.47	20.73	19.37
zambia	16.69	21.72	21.09	19.58
zimbabwe	16.58	21.29	22.49	20.63
total	19.82	23.35	24.52	22.23

Appendix B: Precipitation normals (Sample means)

country	winter	spring	summer	fall
burkinafaso	2.6	15.83	113.78	133.12
cameroon	60.25	101.94	185.08	228.55
egypt	12.81	7.02	2.3	3.51
ethiopia	19.42	49.21	123.71	117.51
ghana	30.87	59.66	112.4	111.74
kenya	88.38	103.02	84.31	59.95
niger	0.75	3.15	64.05	70.55
senegal	2.23	1.05	47.93	112.72
south africa	31.79	54.96	86.38	68.79
zambia	48.26	57.7	108.58	100.67
zimbabwe	7.54	15.4	138.75	89.98
total	25.85	39.83	96.05	102.4

Table 1: Sample of farms

Country	No. of plots	Irrigated plots	Dryland plots
Burkina Faso	1141	59	1082
Cameroon	1013	145	868
Egypt	1030	1030	0
Ethiopia	932	67	865
Ghana	1210	49	1161
Kenya	862	95	767
Niger	1133	52	1081
Senegal	1362	34	1328
South Africa	283	83	200
Zambia	1009	13	996
Zimbabwe	958	123	835
Total	10933	1750	9183

Table 2: Probit model of whether to irrigate

Dependent variable	Irrigated (1/0)
Temperature winter	0.19*** (6.53)
Temperature spring	-0.46*** (17.15)
Temperature summer	0.14*** (3.86)
Temperature fall	0.15*** (3.56)
Precipitation winter	0.01*** (3.23)
Precipitation spring	-0.01*** (3.54)
Precipitation summer	0.005*** (5.07)
Precipitation fall	0.002* (2.45)
Log (mean flow- m3)	0.06*** (8.17)
Chromic cambisols <i>Medium, steep</i>	-1.16* (2.41)
Eutric cambisols <i>Fine, medium</i>	-2.91 (0.59)
Vertic cambisols <i>Fine</i>	1.13** (2.62)
Vertic cambisols <i>Medium, undulating</i>	4.68 (1.88)
Rhodic ferralsols <i>Fine, hilly, steep</i>	1.27 (.61)
Lithosols <i>Coarse, medium, fine, steep</i>	-4.86 (1.49)
Lithosols/Eutric gleysols <i>Hilly</i>	10.58** (3.69)
Chromic luvisols <i>Medium, undulating, hilly</i>	0.57 (1.82)

Table 2 (continued):

Dependent variable	Irrigated (1/0)
Ferric luvisols	1.12***
<i>Coarse, undulating</i>	(7.9)
Gleyic luvisols	1.17***
	(3.82)
Gleyic luvisols	1.33***
<i>Medium, undulating</i>	(5.04)
Gleyic luvisols	-10.47**
<i>Fine, undulating</i>	(2.77)
Orthic luvisols	-2.69
<i>(Medium, hilly)</i>	(1.26)
Dystric nitosols	-7.99*
<i>(Medium, undulating)</i>	(2.00)
Cambic arenosols	-0.79
	(1.18)
Luvic arenosols	-5.41***
	(4.98)
Luvic arenosols	-0.65***
<i>Coarse, undulating</i>	(5.21)
Eutric gleysols	-3.41***
	(6.07)
Eutric gleysols	-0.41*
<i>Coarse, undulating</i>	(2.20)
Calcic yermosols	4.22***
<i>Coarse, medium, undulating, hilly</i>	(4.11)
Chernozems	-0.42**
	(2.60)
Constant	-0.96***
	(3.84)
R2	0.5121
Wald chi2(30)	3967.66
Observations	10880

* p<0.05; ** p<0.01; *** p<0.001

Table 3: Net revenue regressions

Dependent variable	Endogenous	Endogenous	Exogenous	Exogenous
	Net rev/ha (dryland farm)	Net rev/ha (irrigated farm)	Net rev/ha (dryland farm)	Net rev/ha (irrigated farm)
Temperature winter	-120.61* (2.33)	374.41* (2.34)	-124.8103* (2.41)	380.9378* (2.4)
Temperature winter sq	4.58*** (3.39)	-6.79 (1.53)	4.79*** (3.54)	-5.83 (1.31)
Temperature spring	-18.86 (0.2)	-284.49 (1.57)	-19.97 (0.22)	-328.38 (1.91)
Temperature spring sq	-1.88 (0.98)	2.53 (0.6)	-2.02 (1.05)	1.69 (0.4)
Temperature summer	205.86*** (2.95)	1180.58*** (4.28)	212.32** (3.06)	1233.61*** (4.66)
Temperature summer sq	-2.6 (1.95)	-18.27*** (3.81)	-2.73* (2.05)	-18.96*** (4.06)
Temperature fall	-58.81 (1.04)	-1592.37*** (4.11)	-61.66 (1.09)	-1592.27*** (4.09)
Temperature fall sq	0.26*** (0.22)	29.27*** (3.78)	0.41 (0.34)	29.72*** (3.85)
Precipitation winter	-4.37*** (3.71)	12.13*** (1.86)	-4.33*** (3.67)	12.13 (1.86)
Precipitation winter sq	0.03*** (4.54)	0.003 (0.08)	.03*** (4.61)	0.005 (0.11)
Precipitation spring	4.09*** (3.64)	-12.59* (2.18)	4.01*** (3.56)	-13.1* (2.26)
Precipitation spring sq	-0.01 (1.43)	0.01 (0.13)	-0.01 (1.45)	0.01 (0.17)
Precipitation summer	4.56*** (6.47)	21.65*** (4.45)	4.67*** (6.6)	22.88*** (4.9)
Precipitation summer sq	-0.02*** (5.35)	-0.09*** (4.84)	-.02*** (5.35)	-.09*** (4.91)
Precipitation fall	-1.24 (1.94)	-21.36*** (4.58)	-1.28* (2.01)	-22.18*** (4.82)
Precipitation fall sq	0.01*** (5.01)	0.09*** (5.54)	.01*** (5.02)	.09*** (5.62)

Table 3 (continued):

Dependent variable	Endogenous	Endogenous	Exogenous	Exogenous
	Net rev/ha (dryland farm)	Net rev/ha (irrigated farm)	Net rev/ha (dryland farm)	Net rev/ha (irrigated farm)
Chromic cambisols	-411.09**	-407.91	-439.31***	-559.54
<i>Medium, steep</i>	(3.12)	(0.45)	(3.34)	(0.63)
Eutric cambisols	5245.34*	17791.66	5130.45*	16604.16
<i>Fine, medium</i>	(2.35)	(1.6)	(2.31)	(1.5)
Vertic cambisols	-103.68	839.82***	-85.62	1053.52***
<i>Fine</i>	(1.58)	(3.75)	(1.3)	(5.68)
Rhodic ferralsols	-1868.95**	-330.48	-1857.75*	-62.59
<i>Fine, hilly, steep</i>	(2.99)	(0.08)	(2.98)	(0.02)
Lithosols	545.78	16032.28***	438.83	15186.80***
<i>Coarse, medium, fine, steep</i>	(0.48)	(9.61)	(0.39)	(10.04)
Chromic luvisols	-670.35***	3023.22*	-662.76***	3069.53*
<i>Medium, undulating, hilly</i>	(6.63)	(2.51)	(6.54)	(2.56)
Ferric luvisols	-108.7***	-112.59	-81.45***	91.64
<i>Coarse, undulating</i>	(5.55)	(0.62)	(4.52)	(0.73)
Gleyic luvisols	-181.87***	-788.35*	-147.84***	-382.29*
	(4.38)	(2.56)	(3.62)	(2.52)
Gleyic luvisols	655.41***	-289.07	681.85705***	-46.39
<i>Medium, undulating</i>	(4.43)	(1.2)	(4.61)	(0.25)
Gleyic luvisols	1010.34**	-596.26	997.31532**	-66.85736
<i>Fine, undulating</i>	(3.25)	(1.35)	(3.27)	(0.25)
Orthic luvisols	-791.89	-3745.32*	-844.16545	-4296.4641**
<i>(Medium, hilly)</i>	(1.26)	(2.17)	(1.35)	(2.64)
Chernozems	177.49***	-219.62	183.69602***	-149.10838
	(4.32)	(0.69)	(4.48)	(0.48)
Household electrified (1/0)	119.4***	302.13***	120.62***	304.11***
	(7.44)	(3.43)	(7.55)	(3.48)
Log (household size)	29.81**	124.85*	30.74**	116.90
	(2.72)	(2.03)	(2.81)	(1.91)

Table 3 (continued):

Dependent variable	Endogenous	Endogenous	Exogenous	Exogenous
	Net rev/ha (dryland farm)	Net rev/ha (irrigated farm)	Net rev/ha (dryland farm)	Net rev/ha (irrigated farm)
Area of plot	-0.6*** (3.95)	-0.06* (2.28)	-.61*** (4.00)	-.06* (2.14)
Area of plot sq	.0000965** (2.79)	7.237e-07* (2.2)	.000097** (2.81)	7.278e-07* (2.08)
Log (elevation - meters)	-16.85* (2.31)	63.87 (1.89)	-15.64* (2.15)	62.85 (1.86)
Inverse Mills ratio	-19.22*** (4.82)	-256.95 (1.53)		
Constant	-106.24 (0.18)	4169.51* (2.07)	-143.72 (0.24)	3566.84 (1.75)
R2	0.16	0.26	0.2605	0.16
F	25.58	165.59	227.62	24.79
Observations	9131	1749	1749	9131

* p<0.05; ** p<0.01; *** p<0.001

Soil texture

Coarse: sands, loamy sands and sandy loams with less than 18% clay and more than 65% sand.

Medium: sandy loams, loams, sandy clay loams, silt loams, silt, silty clay loams and clay loams with less than 35% clay and less than 65% sand. The sand fraction may be as high as 82% if a minimum of 18% clay is present.

Fine: clay, silty clays, sandy clays, clay loams, with more than 35% clay.

Soil slope (three slope classes)

Undulating: level to gently undulating, with generally less than 8% slope.

Hilly: rolling to hilly with slopes between 8% and 30%.

Steep: steeply dissected to mountainous, with more than 30% slope.

Table 4a: Seasonal and annual marginal effects, evaluated at the sample means for irrigated farms

Irrigated farms: Marginal temperature coefficients

(Net revenue/°C)

Season	Coefficient	T-stat	95% Confidence intervals (Lower, Upper)	
Winter temperature	173.8	2.00	3.1	344.5
Spring temperature	-198.3	-1.98	-395.3	-1.4
Summer temperature	308.7	3.13	115.5	502.0
Fall temperature	-266.8	-2.37	-487.3	-46.4
Annual temperature	17.3	1.05	-15.0	49.8

Irrigated farms: Marginal precipitation coefficients

(Net revenue/mm/mo)

Season	Coefficient	T-stat	95% Confidence intervals (Lower, Upper)	
Winter precipitation	12.3	2.28	1.7	22.8
Spring precipitation	-12.3	-2.85	-20.7	-3.8
Summer precipitation	12.9	3.86	6.3	19.4
Fall precipitation	-13.1	-3.84	-19.7	-6.4
Annual precipitation	-0.2	0.96	-9.9	9.4

Table 4b: Seasonal and annual marginal effects, evaluated at the sample means for dryland farms

Dryland farms: Marginal temperature effects

(Net revenue/°C)

Season	Coefficient	T-stat	95% Confidence intervals (Lower, Upper)	
Winter temperature	70.9	6.17	48.4	93.4
Spring temperature	-111.5	-8.70	-136.6	-86.3
Summer temperature	76.3	6.89	54.6	98.1
Fall temperature	-47.1	-3.99	-70.2	-24.0
Annual temperature	-11.3	-3.97	-16.9	-5.7

Dryland farms: Marginal precipitation effects

Net revenue/mm/mo

Season	Coefficient	T-stat	95% Confidence intervals (Lower, Upper)	
Winter precipitation	-2.88	-3.29	-4.59	-1.16
Spring precipitation	3.44	4.74	2.02	4.87
Summer precipitation	0.84	3.29	0.34	1.35
Fall precipitation	1.21	3.78	0.58	1.83
Annual precipitation	2.62	5.58	1.70	3.54

Table 5: Marginal annual climate impacts

	Dryland	Irrigated
Mean net revenue (US\$/ha)	325.7	1283.8
Mean annual temperature (°C)	23.2	19.6
Mean annual precipitation (mm/mo)	34.4	71.6
Marginal temperature effect (\$/ha/°C)	-11.34	17.37
Marginal precipitation effect (\$/ha/mm/mo)	2.62	-0.22
Annual temperature elasticity	-0.81*	0.31
Annual precipitation elasticity	0.28*	-0.01

* significant at 5%

Note: Marginal effects and elasticities are evaluated at the mean climate of the dryland and irrigated sample.

Table 6: Comparison of irrigation and welfare estimates for endogenous and exogenous models across different climate scenarios

	Climate scenario	Endogenous approach	Exogenous approach
Mean probability of irrigation	2.5°C Δ in T	0.166	
	5°C Δ in T	0.171	0.162
	-20% Δ in P	0.157	
	+20% Δ in P	0.167	
Δ in welfare	2.5°C Δ in T	-8%	-12%
	5°C Δ in T	-14%	-21%
	-20% Δ in P	-21%	-16%
	+20% Δ in P	18%	14%

Note: The current probability of irrigation is 0.162.




Figure 1a: Relationship between seasonal temperature and the probability of adopting irrigation




Figure 1b: Relationship between annual temperature and the probability of adopting irrigation




Figure 2a: Relationship between seasonal precipitation and the probability of adopting irrigation




Figure 2b: Relationship between annual precipitation and the probability of adopting irrigation




Figure 3: Temperature response functions of irrigated and dryland farms




Figure 4: Precipitation response functions of irrigated and dryland farms